



Plant and Insect Diagnostic Clinic

Iowa State University—2445 ATRB, 2213 Pammel Dr

Ames, Iowa 50011-1101

515-294-0581 hortnews.extension.iastate.edu/pidc

NEMATODE SAMPLE SUBMISSION FORM

Submit samples with forms to address listed above

Billing Contact Information

Please print, (*) are required fields

Name* _____

Signature* _____

Company _____

Address* _____

City, State, Zip* _____

Phone* _____

Email* _____

List any others who we can communicate with and who will receive this report. Provide email.

owner of sample secondary contact

Name: _____

Phone: _____

Email: _____

Service requested for in-state samples. Please check.

DO NOT SEND PAYMENT with your sample. Fees are subject to change.

Visit our website: hortnews.extension.iastate.edu/pidc.

Soybean Cyst Nematode Egg Count

Cost per sample: \$30 in-state/\$35 out-of-state

Complete Nematode Count

Cost per sample: \$40 in-state/\$45 out-of-state

Complete nematode counts are primarily for nematodes that feed on corn or turfgrass. Please contact us about samples from fruits or vegetables for specific sample instructions.

Out-of-state sample: contact the clinic before submitting an out-of-state sample.

FOR OFFICE USE ONLY

Sample number _____

Date received _____

INTERNAL BILLING

Department _____

Worktag _____

Account contact _____

By submitting a sample or image along with this form, you signify that **you have read and agree to our Terms and Conditions** found at hortnews.extension.iastate.edu/pidc/terms_and_conditions.

See our website for sampling instructions. Fees are assessed for insufficient samples.

Fill out all required fields (*) or processing of your sample will be delayed/refused.

ISU Accounts Receivable Office will issue a monthly billing statement. Late fees may be assessed on charges greater than 60 days delinquent and customer shall be responsible for collection costs if account is referred to collection.

County/state:										
Field ID										
County, state (if different for each sample)										
Current crop/plant species										

Lab use only

Clinic identification (lab use only)										
---	--	--	--	--	--	--	--	--	--	--

Nematode Soil Samples

Procedures

Soil samples should NOT be collected when the soil is wet or frozen.
Soybean cyst nematode (SCN) egg count

WHEN: In fall: after harvest and before the soil freezes. In spring: before planting, after the ground has thawed and drained. During growing season: from near stunted and/or yellow soybeans.

HOW:Collect a soil core or 1/4 cup of soil (a subsample) from ten to twenty different locations within an area no larger than 10 to 20 acres using a soil sampling probe, hand trowel, or shovel. Define the sampling areas within a field by agronomic, cropping history, or other logical features (see figure) or divide the field to be sampled into evenly sized areas if conditions are similar throughout the field.

Take care not to sample only from “hot spots” or areas of severely damaged plants. Collect soil from the top 8 inches, directly in the root zone (if in season and soybeans are being grown).

Combine all of the subsamples in a bucket and mix the soil thoroughly. From the mixed soil sample, place approximately 1 to 2 cups of soil into a plastic bag or paper soil test bag.

Nematodes on corn

WHEN: For fields with sandy soils (>70% sand content), soil samples should be collected in the spring, before planting, to check for the needle and sting nematodes. For all soils, if stunting, leaf yellowing, and/or mid-day wilting are observed during the growing season, samples should be collected, but only through the R3 corn growth stage (milk stage).

HOW:Collect 10 or more twelve-inch-deep soil cores from the root zone of unthrifty plants. Do not mix, break up, or otherwise disturb the soil cores. If before the V6 corn growth stage, also collect 5 to 10 root masses from plants; the tops of the plants can be cut off and discarded.


Nematodes on other plants (i.e. vegetables, turf grass)

WHEN: Soil samples can be taken any time of the growing season, but the greatest numbers of nematodes are typically found around the roots of annual plants from mid-summer through fall.

HOW: Collect ten to twenty 6-inch-deep soil cores from the root zone of unthrifty plants. Do not mix, break up, or otherwise disturb the soil cores. Also collect some fibrous or feeder roots from the plants showing damage symptoms.

Shipping Samples

- Place each soil sample in a separate, sealed plastic or paper soil sample bag.
- If roots are being collected, place in a separate plastic bag.
- Use a permanent marker to label each bag with grower's name and either a field name or sample number that corresponds to the information on the front of this form.
- Protect the samples from temperatures above 80 degrees.
- Do not be physically rough with the samples (by dropping or throwing them into a box or cooler, for example).
- Deliver or send the samples for processing as quickly as possible; avoid shipping samples on Thursdays and Fridays so that samples do not sit in delivery trucks over the weekend.



Nematode soil sampling patterns for crop fields with unique features.