

# Ergot Poisoning in Cattle

Along with many other problems that livestock producers have to deal with as a result of the ever changing climate, ergotism can routinely be a concern. Ergot is a fungus that grows on the seed head of cereal grains and grasses. Environmental conditions (cool wet spring followed by hot early summer temperatures) are ideal for the ergot fungus to grow. Delayed harvesting of grass hay because of rain also means that late cut hay may also be at risk of ergotism. Historically rye was commonly affected by the ergot fungus but wheat, barley, oats, brome, fescue, bluegrass, Timothy and other grasses can also be infected. All animals are susceptible to ergot but cattle are often most affected.

The fungus produces toxic compounds called ergot alkaloids. The type and quantity of toxic alkaloids varies depending on the specific fungus infecting the plant, the type of plant, and the environmental conditions. Therefore risk of poisoning and the specific clinical signs may vary depending on climatic conditions and species of fungus present.

To determine if pastures or hay are affected, inspect the seed heads for dark brown, purple or black bodies within the seed head (Figure 1). The fungus produces a mass called the sclerotia that completely replaces the ovary tissue of the plant and is often larger than a typical seed. These sclerotia will eventually fall to the ground and overwinter. There is no practical way to eliminate the ergot-producing fungi from pastures. The incidence and severity of future outbreaks will vary with climatic conditions. Determine presence and severity each year by careful field inspection. Fescue grass may be at additional risk since the toxins produced by endophyte infected fescue are very similar to ergot toxins.

## Clinical Signs

The most common sequela of ergot poisoning is associated with vasoconstriction of the small arteries. Vasoconstriction (constriction of blood vessels) can prevent thermoregulation and result in “summer slump”

during the hot summer. Cattle will commonly develop a rough hair coat, lose weight and have extended periods of time standing in water or shade if available. Gangrenous ergotism (synonymous with fescue foot) is also a result of vasoconstriction in the legs and tail.




Figure 1. Ergot bodies in smooth brome seed head.

Gangrenous ergotism is often associated with cold temperatures but can be seen in the summer also. Initially cattle will be lame usually in the hind limbs first. Swelling at the coronary band develops and the animal will eventually slough its hoof if not removed from the ergot alkaloid in time (Figure 2). Necrosis of the tail and ears can also occur.

As with fescue toxicosis, agalactia (decreased or no milk production) can occur. Finally, nervous ergotism is sometimes reported. Nervous signs may be associated



Figure 2. Gangrenous ergotism at coronary band.  
Photo courtesy of Dr. Steve Ensley.

with vasoconstriction in the brain or other alkaloids that specifically cause hyperexcitability and tremors, especially with forced movement.

## Diagnosis

Lameness associated with ergot poisoning should be differentiated from footrot, fescue toxicosis, frostbite, trauma or vesicular disease. Inspection of seed heads should reveal sclerotia if ergotism is the problem. Samples of grass or hay can be analyzed for the concentration of the ergopeptides. A one pound, randomly-collected sample from the pasture or hay that includes seed heads is adequate for analysis. There is no specific test to detect ergopeptides in cattle.

## Treatment

Treatment of ergotism is primarily focused on removing the animal from the source of the ergot toxins as there is no antidote. Animals can recover if they are removed early enough before severe clinical signs are present. However, once gangrene has begun there is little that can be done.

## Prevention

Grass in pasture or hay should be monitored regularly to determine if the ergot sclerotia is present. Many animals will go off feed or seek other grass that is not infected with the ergot fungus. However, cattle are known to selectively graze the seed head and are more at risk than other animals on pastures. Cattle should be removed from pastures that have ergot sclerotia present. Since the ergot fungus is only in the seed head, grazing of infected pasture before the seed head develops is advised. Grass that has developed a seed head can be clipped before allowing cattle to graze. Hay produced from ergot infested grass may be toxic as well. Often the sclerotia will drop off of the grass as it is being handled, but hay should be inspected to make sure that it does not contain sclerotia.

## Suggested References

- Oswieiler GD. 1993. Ergotism. In *Current Veterinary Therapy 3 Food Animal Practice*, edited by J.L. Howard, pp. 334-336. Philadelphia: WB Saunders.
- Evans TJ, Rottinghaus GE, Casteel SW. 2004. Ergot. In *Clinical Veterinary Toxicology*. Edited by K Plumlee, pp. 239-243 . St Louis, Mosby

---

## Authors

Prepared by G. Dewell, DVM, MS, PhD, Beef Extension Veterinarian and Steve Ensley, DVM, PhD, Veterinary Toxicologist, Iowa State University, Veterinary Diagnostic and Production Animal Medicine.



This publication was peer-reviewed by two independent reviewers using a double-blind process.

## ...and justice for all

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, age, disability, and where applicable, sex, marital status, familial status, parental status, religion, sexual orientation, genetic information, political beliefs, reprisal, or because all or part of an individual's income is derived from any public assistance program. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at 202-720-2600 (voice and TDD). To file a complaint of discrimination, write to USDA, Director, Office of Civil Rights, 1400 Independence Avenue SW, Washington, DC 20250-9410, or call 800-795-3272 (voice) or 202-720-6382 (TDD). USDA is an equal opportunity provider and employer.

Issued in furtherance of Cooperative Extension work, Acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture. Cathann A. Kress, director, Cooperative Extension Service, Iowa State University of Science and Technology, Ames, Iowa.